

CERTIFICATE OF VERIFICATION

I, Masayuki HARABAYASHI,
of c/o Morita & Associates, 5th Fl., Itabashi-Chuo Bldg.,
67-8, Itabashi 2-chome, Itabashi-ku, Tokyo 173-0004, Japan,
state that the attached document is a true and complete
translation to the best of my knowledge of Japanese Patent
Application No. 2003-076254.

Dated this 24th day of January, 2008

Signature of translator: M. Harabayashi

[DOCUMENT NAME] Patent Application
[REFERENCE NUMBER] TAK02001P
[ADDRESSEE] Director-General of Patent Office
[INTERNATIONAL PATENT CLASSIFICATION] C12N 15/00
[INVENTOR]

[Address or Residence]
6-8, Narashinodai 1-chome, Funabashi-shi,
Chiba 274-0063, Japan

[Name] Hiroshi TAKAKU

[INVENTOR]

[Address or Residence]
202, 2-1, Ojima 1-chome, Koto-ku,
Tokyo 136-0072, Japan

[Name] Naoko KUROSAKI

[APPLICANT]

[Identification No.] 591028577
[Name] JUNSEI CHEMICAL CO., LTD.

[ATTORNEY]

[Identification No.] 100090251
[Patent Attorney]
[Name] Kenichi MORITA

[INDICATION OF FEE]

[Deposit Account Number] 017813
[Fee] 21000

[LIST OF SUBMITTED DOCUMENTS]

[Name of Document] Specification 1
[Name of Document] Drawings 1
[Name of Document] Abstract 1

[REQUEST FOR A PROOF] Yes

[DOCUMENT NAME] SPECIFICATION

[TITLE OF THE INVENTIONS] NOVEL ANTISENSE OLIGONUCLEOTIDE
AND ANTI-HIV AGENT

[CLAIMS]

[Claim 1] An oligonucleotide consisting of a nucleotide sequence complementary to a nucleotide sequence consisting of at least 15 successive nucleotides in the nucleotide sequence consisting of nucleotides 6-44 of SEQ ID NO: 1.

[Claim 2] An oligonucleotide comprising a nucleotide sequence which specifically hybridizes to a nucleotide sequence consisting of at least 15 successive nucleotides in the nucleotide sequence consisting of nucleotides 6-44 of SEQ ID NO: 1.

[Claim 3] The oligonucleotide according to claim 1 or 2, wherein at least one internucleotide bond is a phosphorothioate bond.

[Claim 4] An anti-HIV agent comprising as an active ingredient the oligonucleotide according to any one of claims 1 to 3.

[DETAILED DESCRIPTION OF THE INVENTION]

[0001]

[Technical Field to which the Invention Pertains]

The present invention relates to a novel antisense oligonucleotide and an anti-HIV agent (an agent for treating and/or preventing HIV).

[0002]

[Prior Art]

As an antiviral agent for a human immunodeficiency virus (hereinafter referred to as HIV), an approach on the basis of an antisense method in which a gene is targeted is known. The antisense method is a technique in which an oligonucleotide having a nucleotide sequence complementary to a target gene is used to inhibit a transcription of the target gene, a splicing of an mRNA, and/or a translation to a protein, thereby specifically inhibiting an expression of the viral protein. One of the most important objects in the antisense method is a selection of a site to be targeted.

[0003]

As antisense oligonucleotides useful as an active ingredient of an anti-HIV agent, for example, oligonucleotides comprising a nucleotide sequence complementary to each nucleotide sequence of a CXCR4 gene or a CCR5 gene [Japanese Unexamined Patent Publication (Kokai) No. 11-292795 (Patent reference 1)], or antisense RNAs for an env portion, env and pol portions, or env, pol, and gag portions [Japanese Translation Publication (Kohyo) No. 2001-502884 (Patent reference 2)] are known.

[0004]

Patent reference 1: Japanese Unexamined Patent Publication (Kokai) No. 11-292795

Patent reference 2: Japanese Translation Publication (Kohyo) No. 2001-502884

[0005]

[Problems to be Solved by the Invention]

An object of the present invention is to provide a novel oligonucleotide useful as an active ingredient of an anti-HIV agent capable of effectively treating and/or preventing an HIV infection, and an anti-HIV agent (an agent

for treating and/or preventing HIV) containing the same.

[0006]

[Means for Solving the Problems]

The object may be attained by the present invention, that is, by an oligonucleotide consisting of a nucleotide sequence complementary to a nucleotide sequence consisting of at least 15 successive nucleotides in the nucleotide sequence consisting of nucleotides 6-44 of SEQ ID NO: 1.

The present invention relates to an oligonucleotide comprising a nucleotide sequence which specifically hybridizes to a nucleotide sequence consisting of at least 15 successive nucleotides in the nucleotide sequence consisting of nucleotides 6-44 of SEQ ID NO: 1.

The present invention relates to an anti-HIV agent comprising as an active ingredient the oligonucleotide.

[0007]

The term "HIV" as used herein means a human immunodeficiency virus, including HIV-1 and variants thereof.

[0008]

[Mode for Carrying out the Invention]

Hereinafter the present invention will be explained.

[1] Oligonucleotide of the present invention

The oligonucleotide of the present invention includes (1) an oligonucleotide consisting of a nucleotide sequence complementary to a nucleotide sequence consisting of at least 15 successive nucleotides in the nucleotide sequence consisting of nucleotides 6-44 of SEQ ID NO: 1, and (2) an oligonucleotide comprising a nucleotide sequence which specifically hybridizes to a nucleotide sequence consisting of at least 15 successive nucleotides in the nucleotide sequence consisting of nucleotides 6-44 of SEQ ID NO: 1.

[0009]

The nucleotide sequence of SEQ ID NO: 1 is a nucleotide sequence of a DIS (Dimerization Initiation Site) region and an adjacent region thereof. The DIS region is located in a DLS (Dimer Linkage Structure) region of HIV-1, and corresponds to the nucleotide sequence consisting of nucleotides 9-43 of SEQ ID NO: 1 (see Figures 6 and 7).

[0010]

As shown in Figures 6 and 7, the DLS of HIV-1 is located, downstream of 5' LTR, at a region containing the initiation codon of a gag gene. It was reported, at the beginning of 1995, that the untranslated flanking sequence was important for a replication step of HIV-1 (Muriaux, D. et al., J. Biol. Chem., 270, 8209-16, 1995), and that the DIS region located in the DLS region was an essential site for a replication of HIV-1, on the basis of in vitro experiments (Berkhout, B. and van Wamel, J. L., RNA, 6, 282-95, 2000; and Damgaard, C. K. et al., Nucleic Acids Res., 26, 3667-76, 1998). The DIS region is located upstream of the initiation codon and SD (Splicing Donor Site) of the gag gene, and it is considered that a stem-loop structure is formed.

However, an antisense method in which the DIS region is targeted was not reported, and it was also not reported whether or not such an approach was effective. In antisense methods, it is well-known that, if a target sequence is essential for a replication or growth of a virus, an antisense oligonucleotide for the target sequence is not necessarily effective as an antiviral agent. Therefore, it could not be predicted, even by those skilled in the art, whether or not antisense oligonucleotides for the DIS region, which was considered to be important for an HIV production, are effective.

[0011]

As shown in Examples described below, the present inventors designed five phosphorothioate oligonucleotides (S-ODNs; Anti-692, Anti-694, Anti-703, Anti-713, and Anti-715), as antisense oligonucleotides for the DIS region and an adjacent region thereof (the nucleotide sequence of SEQ ID NO: 1), and examined an activity of inhibiting the HIV-1 production. As a result, it was found that only three antisense oligonucleotides (Anti-694, Anti-703, and Anti-713) exhibited an excellent activity of inhibiting the HIV-1 production, dose-dependent.

In this connection, as other target sites considered to be important for the HIV production, an antisense

oligonucleotide (28AS) which targeted at the gag region, and an antisense oligonucleotide (PPT-AS) which targeted at a polypurine tract region important for a reverse transcription of viral genes were examined, and it was confirmed that the antisense oligonucleotides did not exhibit the activity of inhibiting the HIV-1 production.

The result that only the DIS region was effective as a target, among plural target regions considered to be important for an HIV production, was not predicted before the experiments, and was unexpected. Further, although it was reported that the DIS region was important for the HIV production, the result that Anti-692 and Anti-715 (the antisense oligonucleotides which contained the DIS region as a target sequence) did not inhibit the HIV production was unexpected, from the view point of the importance of the DIS region.

[0012]

In the oligonucleotide of the present invention consisting of a nucleotide sequence complementary to a nucleotide sequence consisting of at least 15 successive nucleotides in the nucleotide sequence consisting of nucleotides 6-44 of SEQ ID NO: 1, the number of nucleotides is not particularly limited, so long as it is 15 nucleotides or more. The nucleotide number that ensures a specific hybridization to a target region is preferably 15 nucleotides or more, more preferably 18 nucleotides or more, most preferably 20 nucleotides or more. The nucleotide number that ensures a membrane penetration is preferably 30 nucleotides or less, more preferably 28 nucleotides or less, most preferably 25 nucleotides or less. The oligonucleotide of the present invention consists of preferably 15 to 30 nucleotides, more preferably 18 to 28 nucleotides, still further preferably 20 to 25 nucleotides, most preferably 20 nucleotides.

[0013]

The oligonucleotide of the present invention comprising a nucleotide sequence which specifically hybridizes to a nucleotide sequence consisting of at least 15 successive nucleotides in the nucleotide sequence consisting of

nucleotides 6-44 of SEQ ID NO: 1 is not particular limited, so long as it can hybridize to the target nucleotide sequence under the same conditions as those in a living body (for example, in a liquid medium at 37°C) and exhibits an anti-HIV activity. The oligonucleotide of the present invention includes, for example, an oligonucleotide exhibiting an anti-HIV activity, and comprising a nucleotide sequence complementary to a nucleotide sequence consisting of at least 15 successive nucleotides in the nucleotide sequence consisting of nucleotides 6-44 of SEQ ID NO: 1; and an oligonucleotide exhibiting an anti-HIV activity, and comprising a nucleotide sequence complementary to a nucleotide sequence in which one or several nucleotides (preferably one or two nucleotides, more preferably one nucleotide) are substituted, inserted, and/or deleted in a nucleotide sequence consisting of at least 15 successive nucleotides in the nucleotide sequence consisting of nucleotides 6-44 of SEQ ID NO: 1.

[0014]

As the oligonucleotide exhibiting an anti-HIV activity, and comprising a nucleotide sequence complementary to a nucleotide sequence consisting of at least 15 successive nucleotides in the nucleotide sequence consisting of nucleotides 6-44 of SEQ ID NO: 1, there may be mentioned, for example, an oligonucleotide exhibiting an anti-HIV activity, and consisting of a nucleotide sequence in which one or several nucleotides (preferably 1 to 10 nucleotides, more preferably 1 to 5 nucleotides, still further preferably 1 to 3 nucleotides, still further preferably 1 or 2 nucleotides, most preferably 1 nucleotide) are added to the 5' terminus and/or the 3' terminus of a nucleotide sequence complementary to a nucleotide sequence consisting of at least 15 successive nucleotides in the nucleotide sequence consisting of nucleotides 6-44 of SEQ ID NO: 1; more particularly, an oligonucleotide consisting of any one of the nucleotide sequences of SEQ ID NOS: 3 to 5 wherein each of the

internucleotide bonds between nucleosides may be independently a phosphodiester bond or a modified phosphodiester bond; or an oligonucleotide exhibiting an anti-HIV activity, and consisting of a nucleotide sequence in which one or several nucleotides (preferably 1 to 10 nucleotides, more preferably 1 to 5 nucleotides, still further preferably 1 to 3 nucleotides, still further preferably 1 or 2 nucleotides, most preferably 1 nucleotide) are added to the 5' terminus and/or the 3' terminus of any one of the nucleotide sequences of SEQ ID NOS: 3 to 5.

In this connection, as the nucleotide(s) added to the 5' terminus and/or the 3' terminus thereof, a nucleotide complementary to a corresponding nucleotide in the nucleotide sequence of SEQ ID NO: 1 is preferable.

[0015]

The oligonucleotide of the present invention may be prepared from deoxyribonucleosides, ribonucleosides, and/or modified ribonucleosides thereof, such as 2'-O-modified ribonucleosides, so long as the resulting oligonucleotide can function as an antisense oligonucleotide. A preferable modified ribonucleoside is 2'-O-methylribonucleoside, in view of a strong binding property thereof with the base sequence of the target.

Therefore, the oligonucleotide of the present invention may be an oligoribonucleotide (RNA) composed of ribonucleosides and/or modified ribonucleosides, an oligodeoxyribonucleotide (DNA) composed only of deoxyribonucleosides, or a chimera oligoribo/deoxyribonucleotide (RNA/DNA) composed of ribonucleosides (and/or modified ribonucleosides) and deoxyribonucleosides.

[0016]

In the oligonucleotide of the present invention, internucleotide bonds between nucleosides may be independently a phosphodiester bond or a modified phosphodiester bond. The modified phosphodiester bond may be, for example, a methylphosphonate bond wherein one of two non-crosslinked oxygen atoms in the phosphodiester bond is

replaced with a methyl group; a phosphoroamidate bond wherein one of two non-crosslinked oxygen atoms in the phosphodiester bond is replaced with an amino group or a substituted amino group; or a phosphorothioate bond wherein one of two non-crosslinked oxygen atoms in the phosphodiester bond is replaced with a sulfur atom. The oligonucleotide may contain one or more modified phosphodiester bonds as above in one or more internucleotide bonds between nucleosides.

[0017]

The phosphodiester bond is preferable, from the standpoints of the specificity to a nucleotide sequence, an easy procedure for preparation, and a cost of production, and the modified phosphodiester bond is preferable, from the standpoints of a stability of the double-stranded chain, a resistance to a nuclease, a penetrating property through a cell membrane, a low cytotoxicity, and a moderate metabolizability. Further, the phosphorothioate bond is more preferable from the standpoint of stability in a living body.

[0018]

The oligonucleotide used in the present invention may be prepared by known methods. For example, the oligonucleotide may be prepared by an automated DNA/RNA synthesizer in accordance with a conventional phosphodiester method or phosphotriester method, such as an H-phosphonate method or a phosphoramidite method, except for a site to which a 2'-O-methylribonucleotide or a phosphorothioate bond is introduced.

[0019]

The oligonucleotide having phosphorothioate bonds may be prepared, for example, using a 15% N,N,N',N'-tetraethylthiurumdisulfide/acetonitrile solution instead of a water/iodine/pyridine solution that is an oxidizing agent used in a conventional synthesis of polynucleotide.

[0020]

The oligonucleotide having 2'-O-methylribonucleotides may be prepared, for example, by an automated DNA/RNA synthesizer in accordance with the phosphoramidite method,

using a 5'-dimethoxytrityl-2'-O-methylribonucleoside-3'-[(2-cyanoethyl)-(N,N-diisopropyl)]-phosphoroamidite unit.

[0021]

[2] Anti-HIV agent of the present invention

The oligonucleotide of the present invention exhibits an anti-HIV activity, and is useful as an active ingredient of the anti-HIV agent according to the present invention. The term "anti-HIV activity" as used herein means, but is by no means limited to, for example, an activity of inhibiting a production of HIV (particularly HIV-1), or an activity of suppressing an expression of mRNAs or proteins of HIV (particularly HIV-1).

[0022]

The anti-HIV agent of the present invention may contain as the active ingredient the oligonucleotide of the present invention alone, or optionally with a pharmaceutically or veterinarily acceptable known carrier or diluent, and may be prepared in accordance with known formulating techniques (for example, JP 11-292795) of pharmaceutical compositions containing an antisense oligonucleotide.

As the anti-HIV agent of the present invention, there may be mentioned, for example, an anti-HIV agent containing the oligonucleotide of the present invention, an anti-HIV agent containing the oligonucleotide of the present invention and a liposome stable in blood, or an anti-HIV agent containing a vector comprising the oligonucleotide of the present invention.

[0023]

The anti-HIV agent of the present invention may be administered via any of an oral, parenteral or local route. The dose may vary with the species of the subject to be treated (a mammal, particularly a human), a response of the subject to the medicine, a formulation of the medicine, an administration time, an interval of administrations, or the like, but may generally be about 500 mg to about 5000 mg/day.

[0024]

The anti-HIV agent of the present invention may be administered in the form of the oligonucleotide of the present invention, the liposome stable in blood, or the

vector comprising the oligonucleotide of the present invention, and optionally, with a pharmaceutically acceptable known carrier or dilute via any of the oral, parenteral or local routes, once or a multiple of times. The anti-HIV agent of the present invention may be variously formulated to produce, for example, tablets, capsules, lozenges, troches, hard candies, powders, sprays, creams, ointments, suppositories, jellies, gels, pastes, lotions, salves, aqueous suspensions, solutions for injection, elixirs, syrups, or the like.

[0025]

[EXAMPLES]

The present invention now will be further illustrated by, but is by no means limited to, the following Examples.

Example 1: Preparation of antisense oligonucleotides

In this example, three antisense oligonucleotides and two random oligonucleotides were synthesized on the basis of the nucleotide sequence of the DIS (Dimerization Initiation Site) located in the DLS (Dimer Linkage Structure) region of HIV-1, as shown in Figure 1. All oligonucleotides were DNAs in which all internucleotide bonds were phosphorothioate bonds, i.e., phosphorothioate oligonucleotides (S-ODNs).

In Figure 1, each of the numbers in the nucleotide sequence of SEQ ID NO: 1, such as "690" or "700", are nucleotide numbers, when the U3 region in the 5' LTR of a HIV-1 NL432 strain is regarded as the starting point. The terms "Stem", "Loop", and "Bulge" mean a stem region, a loop region, and a bulge region, respectively.

[0026]

An oligonucleotide "Anti-703" (SEQ ID NO: 4) was synthesized as the oligonucleotide of the present invention. Oligonucleotides "Anti-692" (SEQ ID NO: 2) and "Anti-715" (SEQ ID NO: 6) were synthesized as oligonucleotides for comparison. Oligonucleotides "Random" (SEQ ID NO: 7) and "703-Scramble" (SEQ ID NO: 8) were synthesized as control oligonucleotides not having a sequence complementary to the HIV-1 RNA. The oligonucleotide "703-Scramble" shows the same GC content as that of the oligonucleotide "Anti-703" of the present invention.

[0027]

Example 2: Evaluation of S-ODNs for anti-HIV activity using plasmid pNL4-3

(1) Transfection with S-ODNs and plasmid pNL4-3

In this example, a 293T cell (ATCC No. CRL-11268) derived from a human kidney was transfected with an HIV-1 expression vector, plasmid pNL4-3 (Adach, a. et al., J. Virol., 59, 248, 1986), to evaluate S-ODNs prepared in Example 1 with respect to an activity of inhibiting an HIV-1 production.

When the 293T cells are transfected with plasmid pNL4-3, an LTR promoter thereof causes an expression of genes, and HIV-1 is produced. In this example, an amount of a HIV-1 protein, a p24 antigen, produced in the cell and secreted to the culture supernatant was measured.

[0028]

The transfection of the 293T cell with S-ODNs and plasmid pNL4-3 was carried out using a commercially available transfection reagent (FuGENETM6 Transfection Reagent; Boehringer Mannheim, L.L. C, USA) in accordance with a protocol attached thereto.

More particularly, the 293T cells (10^5 cells/2 mL medium/well) were added to each well of a 6-well plate, and incubated in an incubator (at 37°C and 5% CO₂) overnight. As the medium, a Dulbecco's modified Eagle's medium (D-MEM) supplemented with 10% fetal bovine serum (FBS) was used.

[0029]

After the overnight incubation, it was confirmed that the cells adhered to the wells. After the medium was removed from each well, 900 μ L of a serum-free RPMI-1640 medium was added, and the plate was placed in the incubator (at 37°C and 5% CO₂). Further, 100 μ L of each FuGENETM6/S-ODN solution, which had been previously prepared by diluting the transfection reagent (FuGENETM6) and each S-ODN with the serum-free RPMI-1640 medium to predetermined concentrations [S-ODN = 1 μ mol/L, 5 μ mol/L, and 10 μ mol/L (three levels); FuGENETM6 = 1/50 dilution], was added to each well. The plate was gently shaken to a mix in each well, and placed in the incubator (at 37°C and 5% CO₂) for two hours.

After two hours had passed from the addition of the FuGENE™6/S-ODN solution, 110 µL of FuGENE™6/pNL4-3 solution, which had been previously prepared by diluting the transfection reagent (FuGENE™6) and plasmid pNL4-3 with the serum-free RPMI-1640 medium to predetermined concentrations [plasmid pNL4-3 = 0.01 µg/µL; FuGENE™6 = 1/50 dilution], was further added to each well. The plate was gently shaken to a mix in each well, and placed in the incubator (at 37°C and 5% CO₂) for two hours.

After two hours had passed from the addition of the FuGENE™6/pNL4-3 solution, the medium was removed from each well, and each well was washed with 1 mL of phosphate buffered saline [PBS(-)] three times. After 3 mL of an RPMI-1640 medium containing 10% FBS was added to each well, the plate was placed in the incubator (at 37°C and 5% CO₂), and incubated for 48 hours.

[0030]

(2) Purification of HIV-1 and intracellular proteins

After 48 hours from the transfection, 500 µL of each culture supernatant was passed through a 0.45 µm filter, and each filtrate containing purified HIV-1 was transferred to a 1.5 mL tube.

With respect to the cells, each well after removing the supernatant was washed with 1 mL of PBS(-) twice, and 0.5 mL of a 0.05% trypsin-EDTA solution was added to detach the cells from the surface of each well. To each well, 0.5 mL of PBS(-) was added, and the whole was mixed. Each cell suspension was transferred to a 1.5 mL tube, and centrifuged at 2000 rpm for 5 minutes at room temperature to remove the supernatant PBS(-). To the precipitated cells in each tube, 300 µL of a solution for cell lysis (PicaGene; Wako Pure Chemical Industries, Ltd., Osaka, Japan) was added to carry out cell lysis. After incubation at 37°C for 15 minutes, a centrifugation at 13000 rpm was carried out at room temperature for 10 minutes. Each obtained supernatant was transferred to a 1.5 mL tube, and used as a purified solution of intracellular proteins in the following experiments.

[0031]

(3) Measurement of amount of p24 antigen

An amount of an HIV-1 p24 antigen contained in each of the obtained culture supernatants and intracellular proteins was measured by CLEIA (Chemiluminescent Enzyme Immunoassay).

More particularly, each sample was diluted with a commercially available solution for dilution (LUMIPULSE; Fujirebio Inc., Tokyo, Japan) to appropriate concentrations (1/10 to 1/10,000). To 250 μ L of each diluted sample solution, 50 μ L of a commercially available solution for treating an HIV-1 p24 sample (LUMIPULSE I; Fujirebio Inc., Tokyo, Japan) was added, and the whole was mixed gently. A commercially available reagent (LUMIPULSE f; Fujirebio Inc., Tokyo, Japan) was used to measure an amount of the HIV-1 p24 antigen contained therein.

[0032]

The amounts of the HIV-1 p24 antigen contained in the culture supernatants are shown in Figure 2, and the amounts of the HIV-1 p24 antigen contained in the intracellular proteins are shown in Figure 3.

In Figures 2 and 3, the "NL4-3" means a control in which no S-ODNs were added.

As shown in Figure 2, the S-ODN "Anti-703" significantly inhibited the HIV-1 production, and an extremely high activity (99% or more) of inhibiting the HIV-1 production was confirmed. Generally, an antisense nucleic acid needs nucleotides of 17mer or more as a target sequence, and it is theoretically considered that there is no homological sequence other than the target sequence in the range of 17mer or more. However, such an antisense nucleic acid may be actually bind to a nucleotide sequence other than the target gene, in accordance with a homology of several nucleotides, and thus problems such as a high cytotoxicity or an unexpected specificity may be caused by suppressing an expression of genes other than the target. In this experiment, the amount of virus production was inhibited by the S-ODN Anti-703, in accordance with the amounts of Anti-703 used for transfection (i.e., 0.1 μ mol/L, 0.5 μ mol/L, and 1.0 μ mol/L), and the result suggested that the

Anti-703 inhibited the HIV-1 production by specifically binding to the target sequence.

In addition, the decrease in the amount of p24 contained in the supernatant correlated to the decrease in the amount of p24 contained in intracellular proteins shown in Figure 3, and the result shows that the inhibitory activity of Anti-703 for the HIV-1 production inhibits the translation of proteins and the HIV-1 production by specifically recognizing the DIS sequence of HIV-1.

[0033]

Example 3: Evaluation of S-ODNs for anti-HIV activity using plasmid pNL-luc

(1) Transfection with S-ODNs and plasmid pNL4-3

In Example 2, the inhibitory effect of the Anti-703 for the HIV-1 production was shown. Further, it was shown that the inhibitory effect inhibited the production of an HIV-1 protein, p24, by the high antisense activity. As shown in Figure 8, the HIV-1 p24 protein is a structural protein generated from a Gag precursor protein, and the mRNA thereof is translated from the same RNA as the HIV-1 genomic RNA. Since the target DIS region of HIV-1 is located upstream of SD (see Figure 7) and exists in all mRNAs of HIV-1, a plasmid pNL-luc was used to examine whether or not the selected S-ODN acts on other mRNAs efficiently.

[0034]

The plasmid pNL-luc is a plasmid in which a nef region and an env region were deleted and a luciferase gene was inserted instead thereof (see Figure 8). The HIV-1 genomic RNA transcribed in the nucleus was spliced by a splicing mechanism in the nucleus, to become mRNAs capable of expressing proteins. The p24 antigen evaluated in Example 2 is a gene product translated from the HIV-1 genomic RNA, and the luciferase protein is translated from an mRNA generated by the splicing in the nucleus. The plasmid pNL-luc was used to evaluate the advantage of HIV-1 DIS as the target.

[0035]

More particularly, the 293T cells (5×10^5 cells/2 mL medium/well) were added to each well of a 6-well plate, and incubated in an incubator (at 37°C and 5% CO₂). After 14

hours from the addition, it was confirmed that the cells adhered to the wells. The cell density was approximately 30%.

After the medium was removed from each well, 900 μL of a serum-free D-MEM was added, and the plate was placed in the incubator (at 37°C and 5% CO_2). After 15 minutes, 100 μL of each FuGENETM6/S-ODN solution, which had been previously prepared by diluting the transfection reagent (FuGENETM6) and each S-ODN with the serum-free D-MEM to predetermined concentrations [S-ODN = 10 $\mu\text{mol/L}$; FuGENETM6 = 1/50 dilution], was added to each well. The plate was gently shaken to a mix in each well, and placed in the incubator (at 37°C and 5% CO_2) for two hours.

After two hours had passed from the addition of the FuGENETM6/S-ODN solution, 100 μL of FuGENETM6/pNL-luc solution, which had been previously prepared by diluting the transfection reagent (FuGENETM6) and plasmid pNL-luc with the serum-free D-MEM to predetermined concentrations [plasmid pNL-luc = 0.01 $\mu\text{g}/\mu\text{L}$; FuGENETM6 = 1/50 dilution], was further added to each well. The plate was gently shaken to a mix in each well, and placed in the incubator (at 37°C and 5% CO_2) for two hours.

After two hours had passed from the addition of the FuGENETM6/pNL-luc solution, the medium was removed from each well, and 2 mL of D-MEM containing 10% FBS was added to each well. The plate was placed in the incubator (at 37°C and 5% CO_2), and incubated for 48 hours.

[0036]

(2) Purification of intracellular proteins

After 48 hours from the transfection, the culture supernatant was removed from each well. Each well was washed with 1 mL of PBS(-) twice, 0.5 mL of a 0.05% trypsin-EDTA solution was added thereto, and the plate was incubated in the incubator (at 37°C and 5% CO_2) for 5 minutes to detach the cells from the surface of each well. To each well, 0.5 mL of PBS(-) was added, and the whole was mixed. Each cell suspension was transferred to a 1.5 mL tube, and centrifuged at 2000 rpm for 3 minutes at room temperature to remove the supernatant PBS(-). In this connection, 50 μL of

the cell suspension before the centrifugation was used for counting the cell numbers. The precipitated cells in each tube were suspended in 300 µL of PBS(-), and 150 µL of the cell suspension was transferred to another 1.5 mL tube for use in the following Example 3(4). The remaining cell suspension (150 µL) was centrifuged at 2000 rpm for 5 minutes at room temperature to remove the supernatant PBS(-). To the precipitated cells in each tube, 100 µL of a solution for cell lysis (PicaGene; Wako Pure Chemical Industries, Ltd., Osaka, Japan) was added to carry out cell lysis. A centrifugation at 13000 rpm was carried out at room temperature for 10 minutes. Each obtained supernatant was transferred to another 1.5 mL tube, and used as a purified solution of intracellular proteins.

[0037]

(3) Measurement of luciferase activity in intracellular proteins

In this example, 100 µL of a substrate solution for luminescence (Toyo Ink) was added to 10 µL of each purified solution of intracellular proteins, and an luciferase activity in the purified solution of intracellular proteins was measured by a luminometer (LUMAT LB 9507; BERTHOLD).

The result is shown in Figure 4. In Figure 4, "Ran", "703-Scr", and "NL-luc" mean the S-ODN "Random", the S-ODN "703-Scramble", and a control in which no S-ODNs were added, respectively.

[0038]

As shown in Figure 4, when an amount of the HIV-1 gene expressed in cells was evaluated as the luciferase activity, the S-ODN which targeted at the DIS of HIV-1 exhibited the inhibitory effect. Particularly, the Anti-703 significantly inhibited the protein expression in this luciferase activity experiment, as well as the inhibition of the HIV-1 p24 antigen expression. It was known that, as a function of antisense nucleic acids, a binding thereof with the target RNA inhibited progress of ribosomes in translation and suppressed protein synthesis. It is considered that the efficient inhibitory effect of the S-ODN Anti-703 for the HIV-1 production is caused by an effective actions thereof

treated with DNase I, and a reverse transcriptase-polymerase chain reaction (RT-PCR) was carried out. As the primers for RT-PCR, primers capable of specifically amplifying the gag region of HIV-1 RNA were used. As primers for an internal control, primers capable of amplifying a glyceraldehyde-3-phosphate dehydrogenase (G3PDH) gene were used.

[0042]

The result is shown in Figure 5. In Figure 5, "M" means a DNA marker; "NL-luc" means a control in which the plasmid pNL-luc was added, but no S-ODNs were added; "N.C." means a control in which the plasmid pNL-luc was not added, and no S-ODNs were added; and "RT(-)pNL-luc" means a control in which the plasmid pNL-luc was added, but the reverse transcriptase was not added.

As shown in Figure 5, when any one of the S-ODN "Random", the S-ODN "703-Scramble", or the plasmid pNL-luc was transfected singly, no change in the fluorescence of each band was observed. In contrast, a decreased amount of the HIV-1 RNA gene expressed was observed in the S-ODN which targeted the DIS of HIV-1, that is, the RNA gene was not amplified by the RT-PCR in the Anti-703.

From the result, it is considered that the effective inhibitory effect of the Anti-703 for the HIV-1 production is caused by the inhibition of the intracellular protein synthesis and RNA digestion by binding to the target RNA, i.e., by the function of the Anti-703 as antisense nucleic acids.

[0043]

As described above, it was found that the inhibitory effect of the selected S-ODN for the HIV-1 production was caused by the decrease of the HIV-1 protein expression and the decrease of the amount of RNA, and that the S-ODN (Anti-703) specifically bound to the target RNA (DIS in HIV-1) and inhibited the production of viral particles. Further, from the analysis of HIV-1 protein expression, it was confirmed that the DIS region of HIV-1 was important for not only genomic RNA but also mRNAs, and was a effective target site in the antisense method.

[0044]

Example 4: Confirmation of anti-HIV-1 activity using human peripheral blood lymphocytes (1)

In this example, the Anti-703 used in Examples 2 and 3, an Anti-694 (SEQ ID NO: 3), which targeted at a sequence in which the target sequence of the Anti-703 was shifted by 10 nucleotides to the 5' direction, and an Anti-713 (SEQ ID NO: 5), which targeted at a sequence in which the target sequence of the Anti-703 was shifted by 10 nucleotides to the 3' direction, were evaluated using human peripheral blood lymphocytes, with respect to the anti-HIV-1 activity thereof.

Human lymphocytes were separated from peripheral blood taken from a healthy person. An HIV-1 NL432 stain (Adachi, A. et al., J. Virol., 59, 284, 1986) was added to the lymphocytes, and the mixture was allowed to stand on ice for 90 minutes. The lymphocytes were washed with an RPMI-1640 medium to remove any non-adsorbed virus. Each S-ODN (0.5 μ mol/L) such as Anti-703 was mixed with a transfection reagent (DMR1E-C; Invitrogen), and each mixture was added to the infected lymphocytes. The cells were cultivated while the medium was changed every 4 days. After 14 days, an amount of the viral protein HIV-1 p25 antigen produced in the culture supernatant was measured.

[0045]

The result is shown in Figure 9. In Figure 9, "PC" means a positive control in which no S-ODNs were added.

As shown in Figure 9, three S-ODNs other than the S-ODN (Scramble) having a scramble sequence of the Anti-703, i.e., Anti-703, Anti-694, and Anti-713, exhibited an extremely high antiviral activity in comparison with the positive control. The result shows that these S-ODNs can inhibit the viral infection to human peripheral blood lymphocytes.

[0046]

Example 5: Confirmation of anti-HIV-1 activity using human peripheral blood lymphocytes (2)

The procedure described in Example 4 was repeated, except that an S-ODN 28AS (SEQ ID NO: 9) which targeted the gag region, and an S-ODN PPT-AS (SEQ ID NO: 10) which

targeted a polypurine tract region important for a reverse transcription of viral genes were used instead of the Anti-694 and the Anti-713.

The result is shown in Figure 10. In Figure 10, "703AS" means the Anti-703, and "PC" means a positive control in which no S-ODNs were added.

The 28AS which targeted the gag region, or the PPT-AS which targeted the polypurine tract region important for a reverse transcription of viral genes, did not inhibit the viral protein expression, in comparison with the Anti-703 which targeted the DIS region.

[0047]

[Effects of the Invention]

The oligonucleotide of the present invention is useful as an active ingredient of an anti-HIV agent. According to the anti-HIV agent of the present invention, an HIV infection may be effectively treated and/or prevented.

[0048]

[FREE TEXT IN SEQUENCE LISTING]

In each of nucleotide sequences of SEQ ID NOS: 2 to 10, all internucleotide bonds are phosphorothioate bonds.

Each of the nucleotide sequences of SEQ ID NOS: 7 and 8 is an artificially synthesized random sequence.

[0049]

[Sequence Listing]

<110> JUNSEI CHEMICAL CO., LTD.

<120> Novel antisense oligonucleotide and anti-HIV agent

<130> TAK02001P

<160> 10

<210> 1

<211> 49

<212> RNA

<213> Human immunodeficiency virus type 1

<400> 1
acgcaggacu cggcuugcug aagcgcgcac ggcaaggagc gaggggcgg 49

<210> 2
<211> 20
<212> DNA
<213> Human immunodeficiency virus type 1

<220>
<221> misc_feature
<222> (1)..(20)
<223> All internucleotide bonds are phosphorothioate bonds.

<400> 2
cttcagcaag ccgagtcctg 20

<210> 3
<211> 20
<212> DNA
<213> Human immunodeficiency virus type 1

<220>
<221> misc_feature
<222> (1)..(20)
<223> All internucleotide bonds are phosphorothioate bonds.

<400> 3
cgcttcagca agccgagtc 20

<210> 4
<211> 20
<212> DNA
<213> Human immunodeficiency virus type 1

<220>

<221> misc_feature

<222> (1)..(20)

<223> All internucleotide bonds are phosphorothioate bonds.

<400> 4

tgccgtgcgc gcttcagcaa

20

<210> 5

<211> 20

<212> DNA

<213> Human immunodeficiency virus type 1

<220>

<221> misc_feature

<222> (1)..(20)

<223> All internucleotide bonds are phosphorothioate bonds.

<400> 5

cctgcctct tgcggtgcgc

20

<210> 6

<211> 19

<212> DNA

<213> Human immunodeficiency virus type 1

<220>

<221> misc_feature

<222> (1)..(19)

<223> All internucleotide bonds are phosphorothioate bonds.

<400> 6

ccctgcctc ttgcggtgc

19

<210> 7
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: an artificially synthesized
random sequence

<220>
<221> misc_feature
<222> (1)..(20)
<223> All internucleotide bonds are phosphorothioate bonds.

<400> 7
gcgtcatagc tcttgaggac 20

<210> 8
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: an artificially synthesized
random sequence

<220>
<221> misc_feature
<222> (1)..(20)
<223> All internucleotide bonds are phosphorothioate bonds.

<400> 8
cggttagccc atgcgcatgc 20

<210> 9

<211> 28
<212> DNA
<213> Human immunodeficiency virus type 1

<220>
<221> misc_feature
<222> (1)..(20)
<223> All internucleotide bonds are phosphorothioate bonds.

<400> 9
cgctctcgca cccatctctc tcctttta 28

<210> 10
<211> 37
<212> DNA
<213> Human immunodeficiency virus type 1

<220>
<221> misc_feature
<222> (1)..(20)
<223> All internucleotide bonds are phosphorothioate bonds.

<400> 10
ttttcttttg gggggttttt ttcccccott ttctttt 37

[BRIEF DESCRIPTION OF THE DRAWINGS]

[Figure 1]

Figure 1 illustrates the relationship between the nucleotide sequences of S-ODNs used in Examples and the DIS region of HIV-1 as the target sequence.

[Figure 2]

Figure 2 is a graph showing amounts of an HIV-1 p24 antigen in culture supernatants of 293T cells transfected with various concentrations of S-ODNs (0.1 $\mu\text{mol/L}$, 0.5 $\mu\text{mol/L}$, and 1.0 $\mu\text{mol/L}$) and plasmid pNL4-3 (1 μg).

[Figure 3]

Figure 3 is a graph showing amounts of an HIV-1 p24

antigen in intracellular proteins of 293T cells transfected with various concentrations of S-ODNs (0.1 $\mu\text{mol/L}$, 0.5 $\mu\text{mol/L}$, and 1.0 $\mu\text{mol/L}$) and plasmid pNL4-3 (1 μg).

[Figure 4]

Figure 4 is a graph showing a luciferase activity in intracellular proteins of 293T cells transfected with various S-ODNs (1.0 $\mu\text{mol/L}$) and plasmid pNL-luc (1 μg).

[Figure 5]

Figure 5 is a graph showing an expression level of intracellular HIV-1 mRNA gene, detected by RT-PCR, in 293T cells transfected with various S-ODNs (1.0 $\mu\text{mol/L}$) and plasmid pNL-luc (1 μg).

[Figure 6]

Figure 6 schematically illustrates the HIV-1 genomic RNA and the structure of the 5' terminus thereof.

[Figure 7]

Figure 7 schematically illustrates (A) the structure of the 5' terminus of the HIV-1 genomic RNA and (B) the secondary RNA structure model thereof.

[Figure 8]

Figure 8 schematically illustrates the translational mechanisms of (A) HIV-1 genomic RNA and (B) viral mRNA.

[Figure 9]


Figure 9 is a graph showing an antiviral activity with respect to human peripheral blood lymphocytes as the target.

[Figure 10]


Figure 10 is a graph showing an antiviral activity with respect to human peripheral blood lymphocytes as the target.

[DOCUMENT NAME] DRAWINGS


[Figure 1]




[Figure 2]




[Figure 3]




[Figure 4]




[Figure 5]



[Figure 6]




[Figure 7]



[Figure 8]


(A) Genomic RNA




(B) Luciferase protein mRNA



[Figure 9]



[Figure 10]



[DOCUMENT NAME] Abstract

[ABSTRACT]

[OBJECT] A novel antisense oligonucleotide useful as an active ingredient of an anti-HIV agent capable of effectively treating and/or preventing an HIV infection, and an anti-HIV agent (an agent for treating and/or preventing HIV) comprising the same, are disclosed.

[MEANS FOR SOLUTION] The antisense oligonucleotide targets a DIS (Dimerization Initiation Site) region of HIV-1.

[SELECTED DRAWINGS] None